arXiv:1412.2215v1 [astro-ph.HE] 6 Dec 2014

Saha Equation in an Uniformly Accelerated Reference Frame and Some of Its Physical Implications

Sanchari De and Somenath Chakrabarty†
Department of Physics, Visva-Bharati, Santiniketan-731235, India
†somenath.chakrabarty@visva-bharati.ac.in

PACS:03.30.-p,04.20.-q,05.20.Gg

Abstract

The Saha equations for the photo-ionization of hydrogen atoms and the electron positron pair production at high temperature are obtained in a reference frame undergoing a uniform accelerated motion in an otherwise flat Minkowski space-time geometry. Some of the physical implications of our findings are discussed.

1 Introduction

For the space-time coordinate transformation between two inertial frame of references, the well known Lorentz transformations are used [1]. However, from the principle of equivalence, one can treat the space-time transformations in an uniformly accelerated frame in the same manner as it is done in special theory of relativity, i.e., with flat geometrical structure [2, 3, 4]. Let us consider two reference frames- the frame S is unaccelerated inertial one in which there is a uniform gravitational field and the frame S', which is undergoing an accelerated motion with respect to some inertial frame, but there is no gravitational field. Then in accordance with the principle of equivalence these two frames are physically equivalent. Alternatively, one can state, that a reference frame in uniform acceleration in absence of gravitational field is equivalent to a frame at rest in presence of an uniform gravitational field.

Based on the physical concepts of principle of equivalence, in this article first we shall obtain the diagonal elements of metric tensor $g^{\mu\nu}$. Analogous to the Minkowski metric tensor, the off-diagonal terms are also found to be zero. In the next step with the conventional form of action as has been defined in special theory of relativity [1], the Lagrangian of a particle (which may be a hydrogen atom or a hydrogen ion or an electron) is obtained. Which further gives the momentum and energy or Hamiltonian of the particle

from the standard relations. Then considering a partially ionized hydrogen plasma which is an active mixture of neutral hydrogen atoms, hydrogen ions, electrons and photons or considering an electron-positron plasma consisting of an interacting mixture of electron, positron and photons, we shall establish the modified form of Saha equations for both the cases as observed from an uniformly accelerated frame of reference. To the best of our knowledge, the study of Saha equation in uniformly accelerated frame has not been reported earlier. We shall compare our findings with the conventional results.

The article is organized in the following manner: In the next section we shall establish some of the useful relations of special theory of relativity in uniformly accelerated frame. In section-3 we shall investigate photo-ionization of hydrogen atoms in partially ionized hydrogen plasma. In section-4 we shall study the electron-positron pair production from intense electromagnetic radiation at high temperature. In the last section we shall conclude our findings and discuss the physical implication of our results.

2 Basic Formalism

In this section we shall establish some of the useful formulas of special relativity with uniform accelerated motion (see [5, 6]). We assume that the frame S' has rectilinear motion with velocity v, along x-direction with respect to the frame S. Further the coordinates of an event occured at the point P (say) is indicated by (x, y, z, t) in S-frame and with (x', y', z', t') in the frames S'. The primed and the un-primed coordinates are related by the conventional form of Lorentz transformations and are given by

$$x' = \gamma(x - vt), \quad y' = y, \quad z' = z \text{ and } t' = \gamma\left(t - \frac{vx}{c^2}\right) \text{ with } \gamma = \left(1 - \frac{v^2}{c^2}\right)^{-1/2} \tag{1}$$

is the Lorentz factor. Now for the sake of simplicity, we assume that the acceleration of a particle is also along x-axis in S-frame and is α_x. Then in the primed frame it is given by

$$\alpha'_x = \frac{\alpha_x}{\gamma^3\left(1 - \frac{vu_x}{c^2}\right)^3}, \quad \text{where} \quad \alpha_x = \frac{du_x}{dt} \tag{2}$$

Therefore for this particle having constant acceleration in S'-frame, in which it was instantaneously in coincidence at some past time, we have

$$\alpha'_x = g \text{ (say)} = \gamma^3 \alpha_x \tag{3}$$

Here it has further been assumed that the motion of the particle is also rectilinear and is along x-direction with $u_x = v$. Then with the initial condition that $u_x = 0$ at $t = 0$, we have

$$u_x = \frac{gt}{\left[1 + \left(\frac{gt}{c}\right)^2\right]^{1/2}} \tag{4}$$

Therefore the proper time of the particle is given by

$$\tau = \frac{c}{g} \sinh^{-1}\left(\frac{gt}{c}\right) \tag{5}$$

Hence

$$ct = \frac{c^2}{g} \sinh\left(\frac{g\tau}{c}\right) \tag{6}$$

Then from the expression $u_x = dx/dt$, the velocity of the particle along x-direction and using eqn.(4), we have

$$x = \frac{c^2}{g} \cosh\left(\frac{g\tau}{c}\right) \tag{7}$$

Here eqns.(6) and (7) are the parametric form of the equation for the world line of the particle undergoing constant accelerated motion. Hence the hyperbolic motion of the particle can be represented by the equation

$$x^2 - (ct)^2 = \left(\frac{c^2}{g}\right)^2 \tag{8}$$

To obtain the Rindler coordinates, which are appropriate for a system / observer undergoing constant proper acceleration g in a Minkowski flat spacetime, the scheme as given below has been adopted. Following [5, 6], we have constructed a vector which is tangential with the world line and is proportional to

$$\left(\cosh\left(\frac{g\tau}{c}\right), \sinh\left(\frac{g\tau}{c}\right)\right) \tag{9}$$

These are the parametric form of coordinates of the points moving along the proper time axis of the uniformly accelerated observer. Next consider a vector which is orthogonal to above vector. The parametric form of which is proportional to

$$\left(\sinh\left(\frac{g\tau}{c}\right), \cosh\left(\frac{g\tau}{c}\right)\right) \tag{10}$$

Then the parametric form of the world line of a second observer, which is h distance ahead of the first one as measured by the first observer is given by

$$ct = \left(\frac{c^2}{g} + h\right)\sinh\left(\frac{g\tau}{c}\right) \quad \text{and} \quad x = \left(\frac{c^2}{g} + h\right)\cosh\left(\frac{g\tau}{c}\right) \tag{11}$$

For the second accelerated observer, the hyperbolic motion is described by

$$x^2 - (ct)^2 = \left(\frac{c^2}{g} + h\right)^2 = \left(\frac{c^2}{g}\right)^2\left(1 + \frac{gh}{c^2}\right)^2 \tag{12}$$

The magnitude of constant acceleration for the second observer is therefore

$$g' = \frac{g}{\left(1 + \frac{gh}{c^2}\right)} \tag{13}$$

i.e., the two observers undergoing accelerated motions with the different magnitudes of acceleration.

To obtain the Rindler coordinates, we consider an event with space time coordinate ct' and x' as measured by the first observer. Further it has been assumed that x' is the distance from the origin of the first observer and t' is the corresponding proper-time. Then with respect to an inertial frame, we have from the eqns.(8), (9) and (11) and considering simultaneity of these events

$$ct = \left(\frac{c^2}{g} + x'\right)\sinh\left(\frac{gt'}{c}\right) \quad \text{and} \quad x = \left(\frac{c^2}{g} + x'\right)\cosh\left(\frac{gt'}{c}\right) \tag{14}$$

Hence one can also express the inverse relations

$$ct' = \frac{c^2}{2g}\ln\left(\frac{x + ct}{x - ct}\right) \quad \text{and} \quad x' = (x^2 - (ct)^2)^{1/2} - \frac{c^2}{g} \tag{15}$$

The Rindler space-time coordinates, given by eqns.(14) and (15) are then just an accelerated frame transformation of the Minkowski metric of special

relativity. The Rindler coordinate transform the Minkowski line element

$$ds^2 = d(ct)^2 - dx^2 - dy^2 - dz^2 \;\text{ to }\; ds^2 = \left(1 + \frac{gx'}{c^2}\right)^2 d(ct')^2 - dx'^2 - dy'^2 - dz'^2 \tag{16}$$

Since the motion is assumed to be rectilinear and along x-direction, $dy' = dy$ and $dz' = dz$. The general form of metric tensor may then be written as

$$g^{\mu\nu} = \text{diag}\left(\left(1 + \frac{gx}{c^2}\right)^2, -1, -1, -1\right) \tag{17}$$

Now following the concept of relativistic dynamics of special theory of relativity [1], the action integral may be written as (see also [7])

$$S = -\alpha \int_a^b ds \equiv \int_a^b L dt \tag{18}$$

Then using eqns.(16) and (18) and putting $\alpha = -m_0 c$, where m_0 is the rest mass of the particle, the Lagrangian of the particle is given by

$$L = -m_0 c^2 \left[\left(1 + \frac{gx}{c^2}\right)^2 = \frac{v^2}{c^2}\right) \tag{19}$$

where $\vec{v}$ is the velocity of the particle. The momentum of the particle is then given by

$$\vec{p} = \frac{m_0 \vec{v}}{\left[\left(1 + \frac{gx}{c^2}\right)^2 - \frac{v^2}{c^2}\right]^{1/2}} \tag{20}$$

Hence the Hamiltonian of the particle or the single particle energy is given by

$$H = \varepsilon_p = m_0 c^2 \left(1 + \frac{gx}{c^2}\right)\left(1 + \frac{p^2}{m_0^2 c^2}\right)^{1/2} \tag{21}$$

In the forth coming sections, using the expression for single particle energy as given in eqn.(21), we shall obtain the Saha equations for photo-ionization of hydrogen atoms and also for electron-positron pair creation. We shall follow the standard formalism as discussed in the text books on astrophysics and statistical mechanics [9, 8].

3 Photo-Ionization of Hydrogen Atoms

To study the Saha ionization process for hydrogen atoms, we consider a partially ionized hydrogen plasma, consisting of neutral hydrogen atoms, hydrogen ions, electrons and electromagnetic radiation or photons. The system is assumed to be of cylindrical in nature and the plasma is expanding with a constant acceleration g along the positive x-direction, which is the symmetry axis of the cylinder. Then according to principle of equivalence, any ionization or de-ionization of the accelerated particles taking place at some point is equivalent to their occurrence at rest frame in presence of a gravitational field g. Now depending on the amount of energy transferred, the interaction with photon will either excite the hydrogen atoms or ionize them. The electrons can also be captured by the hydrogen ions, which is the de-ionization process. Ultimately a dynamic dead lock situation will arise, when the rate of ionization process and the rate of de-ionization process become just equal, which may be represented by the reaction equation

$$H_n + \gamma \leftrightarrow H^+ + e^- \tag{22}$$

where H_n indicates the nth excited state of hydrogen atom and $n = 1$ is the ground state. Under such physical condition the partially ionized hydrogen plasma will be in thermodynamic equilibrium with all of its constituents. In chemical equilibrium condition the chemical potentials of the components are related by the equation

$$\mu(H_n) = \mu(H^+) + \mu(e) \tag{23}$$

Since the number of photons is not conserved, $\mu(\gamma) = 0$.

In the non-relativistic situation, the single particle energy as shown in eqn.(21) may be approximated as

$$\varepsilon_p = m_0 c^2 \left(1 + \frac{gx}{c^2}\right) + \left(1 + \frac{gx}{c^2}\right) \frac{p^2}{2m_0} \tag{24}$$

Then defining

$$m' = \frac{m_0}{1 + \frac{gx}{c^2}} \quad \text{and} \quad m'' = \left(1 + \frac{gx}{c^2}\right) m_0 \tag{25}$$

Hence we can write

$$\varepsilon_p = m'' c^2 + \frac{p^2}{2m'}$$

as the effective single particle energy or Hamiltonian. Then it can very easily be shown following the standard results of text books on statistical mechanics [10], the number density of a particular component (except for γ)

$$n = \frac{N}{\Delta V} = \frac{4\pi g_d}{h^3} \int_0^\infty p^2 dp \exp\left[-\frac{1}{kT}\left(\frac{p^2}{2m'} + m''c^2 - \mu\right)\right] \tag{26}$$

where g_d is the degeneracy of the species, $\Delta V = A\Delta x$ is a small volume element, Δx is a small length element in the x-direction at a distance x from the origin and A is the cross-sectional area of the cylinder. The length Δx is such that g is constant within ΔV. On evaluating the above integral, we have

$$n = n_Q g_d \exp\left[\frac{1}{kT}\left(\mu - m''c^2\right)\right] \tag{27}$$

Hence

$$\mu = m''c^2 - kT \ln\left(\frac{g_d n_Q}{n}\right) \tag{28}$$

This is the general expression for chemical potential for a particular component in terms of its concentration. Further,

$$n_Q = \left(\frac{2\pi m' kT}{h^2}\right)^{3/2} \tag{29}$$

is called the quantum concentration for that particular species in the mixture. Then using eqn.(23) for chemical equilibrium condition and eqn.(28), the expression for the chemical potential, we have

$$\frac{n(H_n)}{n(H^+)n(e)} = \frac{g_n}{n_{Q_e}} \exp\left(-\frac{\Delta E_n}{kT}\right) = R_{g>0} \text{ (say)} \tag{30}$$

where $n(i)$ are the number density for the species i and $g_n = n^2$ is the degeneracy of the neutral hydrogen atom in the nth excited state (since the degeneracy of electrons has been considered separately in the derivation, the factor 2 has not been there in the expression for g_n) and

$$\Delta E_n = \Delta m'' c^2 = \left(1 + \frac{gx}{c^2}\right) \Delta m c^2, \quad \text{with} \quad \Delta m = m(H_n) - m(H^+) - m(e) \tag{31}$$

is a measure of excitation energy.

Here eqn.(30) gives the ratio of equilibrium concentration at a given temperature and in this particular situation for a given value of $g > 0$. Now the

conventional form of Saha equation, i.e., in an inertial frame with $g = 0$ is given by

$$\frac{n(H_n)}{n(H^+)n(e)}_{g=0} = \frac{g_n}{n_{Q_e\, g=0}} \exp\left(-\frac{\Delta m c^2}{kT}\right) = R_{g=0} \text{ (say)} \tag{32}$$

Hence the ratio with $g > 0$ and $g = 0$ can be written as

$$\frac{R_{g>0}}{R_{g=0}} = \left(1 + \frac{gx}{c^2}\right)^{-3/2} \exp\left(-\frac{gx\Delta m}{kT}\right) \tag{33}$$

It is quite obvious from eqns.(30) and (33) that both the ratios decrease with g. The variation with g is much faster than the fall up of screened coulomb or Yukawa potential.

4 Pair Production at High Temperature

We next consider the electron-positron pair creation at high temperature. The temperature is assumed to be such that kT is larger than at least two times the electron mass. The pair production process can be represented by the reaction equation

$$\gamma + \gamma \leftrightarrow e^- + e^+ \tag{34}$$

Where we have assumed that the interacting electron-positron plasma along with the photons is in thermodynamic equilibrium. In this case the chemical equilibrium condition is given by

$$\mu(e^+) + \mu(e^-) = 0 \tag{35}$$

where

$$\mu(e^-) = m''(e)c^2 - kT \ln\left(\frac{g_e n_Q}{n_{e^-}}\right) \text{ and } \mu(e^+) = m''(e)c^2 - kT \ln\left(\frac{g_e n_Q}{n_{e^+}}\right) \tag{36}$$

Then from eqn.(35), we have

$$n_{e^-} n_{e^+} = g_e^2 n_Q^2 \exp\left(-\frac{2m''(e)c^2}{kT}\right) \tag{37}$$

where

$$n_Q = \left(\frac{2\pi m'(e) kT}{h^2}\right)^{3/2} \tag{38}$$

is the quantum concentration for electron and positron. Here $m'(e)$ and $m''(e)$ are given by eqn.(25) with m_0 replaced by $m(e)$.

If the system is assumed to be charge neutral then $n_{e^-} = n_{e^+}$. Further eqn.(37) can also be expressed as a ratio of the product of electron-positron concentration with $g > 0$ and $g = 0$, and is given by

$$\frac{n_{e^-}n_{e^+\,g>0}}{n_{e^-}n_{e^+\,g=0}} = \left(1 + \frac{gx}{c^2}\right)^{-3/2} \exp\left(-\frac{gxm(e)}{kT}\right) \tag{39}$$

The above equation and the eqn.(37) also show that both the ratios decrease with g in the same manner as that of neutral hydrogen concentration (eqns.(30) and (33)).

5 Conclusion

In section-3 of this article we have shown that the concentration of neutral hydrogen decreases with g. As a consequence it may be observed that the intensity of some of the Lyman α-lines in Lyman α-forest, for which neutral cosmic hydrogen atoms in giant hydrogen clouds are responsible will be low if measured from an uniformly accelerated frame or equivalently, if some of the clouds move with uniform acceleration. The Lyman-α absorption lines in the quasar spectra result from intergalactic neutral hydrogen gas through which the galaxy or quasar's light has traveled. Since neutral hydrogen clouds in the intergalactic medium are at different degrees of redshift (due to their varying distance from Earth), their absorption lines are observed at a range of wavelengths. Each individual cloud leaves its fingerprint as an absorption line at a different position in the observed spectrum. Therefore those Lyman-α, which are produced by the uniformly accelerated neutral hydrogen clouds will be fainter [14]. Our explanation for the less amount of neutral hydrogen in accelerated frame is that an enormous amount of extra energy in the form of intense electromagnetic radiation will be transported to the neutral hydrogen gas cloud from background vacuum states. This electromagnetic radiation causes ionization of more hydrogen atoms and shifts the chemical equilibrium point towards more hydrogen ions. The argument behind such explanation is that an observer in an accelerated frame will see radiation in the vacuum of inertial observer, whereas from inertial frame, there will be no radiation in the vacuum states. Which therefore indicates the vacuum is a relative concept. Therefore our findings are consistent with Unruh effect [12].

The Unruh effect, which is essentially hypothetical in nature, predicts that an accelerating observer will see black-body radiation in a true vacuum for an inertial observer. The temperature of the inertial vacuum as measured by the accelerated observer will increase with the magnitude of acceleration. In other words, the background appears to be warm from an accelerating reference frame. The ground state for an inertial observer is seen as in thermodynamic equilibrium with a non-zero temperature by the uniformly accelerated observer. The Unruh temperature is given by $T_U = \hbar g/(2\pi ck)$ [13]. Whereas the Hawking temperature is given by $T_H = \hbar K_H/(2\pi ck)$ for a Schwarzschild black hole of mass M and $K_H = 1/(4M)$ [11].

In section-4, we have showed that the concentration of electron and positron pairs in the high temperature region filled with an intense electromagnetic radiation decreases with g when observed from an uniformly accelerated frame. In other wards, for an accelerated observer, the creation of electron-positron pairs will be suppressed. The pairs may not be electron-positron, but it can be pairs of any particle and its anti-particle. Even the creation of pairs from single high energy photon in presence of some external field will also be suppressed. The result is again consistent with the prediction that the acclerated observer will not see Hawking radiation [3] (see also [15]).

In this study we have assumed rectilinear motion of S'-frame. However, we can remove this restriction and can assume that the velocity of the primed frame $\vec{v} \equiv (v_x, v_y, v_z)$. One can also assume that the primed frame is moving radially out ward. In the case of a non-rectilinear motion of the acclerated observer, to get Lorentz transformations, the standard technique which generally followed, is to decompose the radial vector $\vec{r}$ into two components, $\vec{r}_{||}$, parallel to the velocity vector $\vec{v}$ and $\vec{r}_{\perp}$ in the orthogonal direction. Under Lorentz transformation, the parallel component will change, whereas the orthogonal component will not be affected. The transformed vector is then given by

$$\vec{r'} = \vec{r} + \frac{\vec{r}.\vec{v}}{v^2}(\gamma - 1)\vec{v} - \gamma t\vec{v} \tag{40}$$

and

$$t' = \gamma\left(t - \frac{\vec{r}.\vec{v}}{c^2}\right) \tag{41}$$

where

$$\vec{r} = \vec{r}_{||} + \vec{r}_{\perp}, \tag{42}$$

$$\vec{r}_{||} = \frac{\vec{r}.\vec{v}}{v^2}\vec{v} \quad \text{and} \quad \vec{r}_{\perp} = \vec{r} - \frac{\vec{r}.\vec{v}}{v^2}\vec{v} \tag{43}$$

Then it is a matter of simple algebra to show that the same one dimensional type results can also be obtained for radially accelerated motion with proper acceleration $\alpha'_r = g$ and from inertial frame $\alpha_r = \alpha'_r\gamma^{-3}$. In this case ΔV is the volume of a thin shell with Δx replaced by Δr, thickness of the shell.

References

[1] L.D. Landau and E.M. Lifshitz, The Classical Theory of Fields, Butterworth-Heimenann, Oxford, 1975.

[2] S. Weinberg, Gravitation and Cosmology, Wiley, New York, 1972; C.W. Misner, Kip S. Thorne and J.A. Wheeler, Gravitation, W.H. freeman and Company, New York, 1972.

[3] W. Rindler, Essential Relativity, Springer-Verlag, New York, 1977

[4] C. Moller, The Theory of Relativity, Calarendon Press, Oxford, 1972.

[5] M. Socolovsky, arXiv.1304.2833.

[6] G.F. Torres del Castillo and C.L. Perez Sanchez, Revista Mexican De Fisika **52**, 70 (2006).

[7] C-G. Huang and J-R. Sun, arXiv:gr-qc/0701078.

[8] L.D. Landau and E.M. Lifshitz, Statistical Physics, Part-I, Butterworth-Heimenann, Oxford, 1980.

[9] A.C. Phillips, The Physics of Stars, John Wiley & Sons, New York, 1996.

[10] K. Huang, Statistical Mechanics, Wiley-Eastern Pvt. Ltd., New Delhi, 1975.

[11] S.L. Shapiro and S.A. Teukolsky, Black Holes, White Dwarfs and Neutron Stars, John Wiley and Sons, New York, (1983).

[12] W.G. Unruh, Phys. Rev. **D14**, 4 (1976); W.G. Unruh, Phys. Rev. **D14**, 870 (1976).

[13] P.M. Alsing and P.W. Milonni, arXiv:quant-ph/0401170.

[14] M. Rauch, Annu. Rev. Astron. Astrophys. **36**, 267 (1998).

[15] S.N. Hawking, Nature, **248**, 30 (1974); S.W. Hawking, Comm. Math. Phys. **43**, 199 (1975).